

ANTICANCER STUDY OF PORCUPINE (*Hystrix
brachyura*) BEZOAR ON MELANOMA AND THE
IDENTIFICATION OF ITS ACTIVE COMPOUNDS

BY

AL'AINA YUHAINIS BT FIRUS KHAN

A thesis submitted in fulfilment of the requirement for the
degree of Doctor of Philosophy in Health Sciences

Kulliyyah of Allied Health Sciences
International Islamic University Malaysia

AUGUST 2020

ABSTRACT

Hystrix brachyura bezoar (PB) has been traditionally used as an alternative medicine to treat cancer. However the anticancer effect of PB is yet to be explored. Therefore this study aims to evaluate PB anticancer potential using metabolomics approach and molecular docking. Four different PBs namely PB-A, PB-B, PB-C and PB-D were procured and used in this study to evaluate the anticancer effect. All PBs were treated with water through ultrasonication-assisted extraction technique to obtain aqueous extracts. The melanoma cells (A375) were chosen for the evaluation of further anticancer effects. A375 cells were subjected to cell proliferation, colony formation, apoptosis, cell cycle arrest, cell migration, cell invasion assays followed by molecular pathways using real time polymerase chain reaction (qPCR) and *in vivo* antiangiogenesis using zebrafish (*Danio rerio*) larvae. Additionally, all four PBs were also evaluated for their toxicity effect using zebrafish embryos. The anticancer compounds of PB were putatively identified using liquid chromatography–mass spectrometry (LC-MS) and liquid chromatography–mass spectrometry (LC-MS) based metabolomics approaches. All the compounds identified were then docked to BCL-2, cyclin B/CDK1 complex, VEGF and NM23 crystal structures to predict the ligand-protein interaction. Lastly, the study also developed a validated regression model using Fourier transform infra-red (FT-IR) spectroscopy to predict the anticancer activity of new PB extract to ensure its quality as an effective anticancer agent. All PB extracts exhibited significant cytotoxicity on A375 cells. Further analysis revealed that PB-A, PB-C and PB-D had a good inhibitory effect on cell proliferation, colony formation, cell cycle, cell migration, cell invasion, angiogenesis and apoptosis inducer. PB-A, PB-C and PB-D revealed apoptosis through intrinsic pathway, arresting the cell cycle at G2/M phase by downregulating cyclin B and CDK1. PB-A, PB-C and PB-D at molecular level were showed to suppress NM23, E-cadherin, MMP2 and MMP9. The toxicity assessment showed the morphological developmental defects caused by all four PB extracts such as deformed brain section, contorted backbone with deformities in somites and notochord, deformities in soft tissues (yolk sac, pericardial edema and swim bladder). Additionally, the PB extracts were showed to affect the cardiovascular systems via presence of heart edema, downregulating the heart rate and blood flow. The LC₅₀ values at 96 hpf were <100 µg/mL. Chemical profiling analyses of PB extracts through GC-MS and LC-MS based metabolomics approached identified 4-androsten-4-ol-3,17-dione, acetate cholest-7-en-3-ol, gallic acid, isolongifolol, mangiferin and propafenone as the active principles. The docking results predicted the interaction of the active principles with BCL-2, cyclin B/CDK1 complex, VEGF and NM23 majorly via hydrophobic interaction with protein residues. The docking results showed good binding affinity to the crystal structures of BCL-2, cyclin B/CDK1 complex, VEGF and NM23 for 4-androsten-4-ol-3,17-dione, cholest-7-en-3beta-ol,4,4-dimethyl-, acetate and mangiferin, suggesting the roles of these compounds as the potential anticancer agents. Moreover, the study has also generated a validated statistical model to predict the anticancer activity of new PB extracts. Conclusively, the study has revealed the anticancer activity of *Hystrix brachyura* bezoar, active principles, *in vitro* mechanism of action and *in vivo* toxicity effect using zebrafish model for the first time. Results of this study further support the traditional claims for the use of *Hystrix brachyura* bezoar as an anticancer agent in Malaysia.

خلاصة البحث

تم استخدام (PB) *Hystrix brachyura bezoar* بشكل تقليدي كدواء بديل لعلاج السرطان. علماً أنه مع ذلك، لم يتم بعد استكشاف التأثير المضاد للسرطان من PB. ولذلك كانت هذه الدراسة تهدف إلى تقييم إمكانات مكافحة السرطان عن طريق مضاد السرطان PB باستخدام نهج الأيض والالتحام الجزيئي. تم اقتناء أربعة PBs مختلفة وهي PB-A و PB-B و PB-C و PB-D واستخدامها في هذه الدراسة لتقييم التأثير المضاد للسرطان. تمت معالجة جميع مركبات ثنائي الفينيل متعدد البروم بالماء من خلال تقنية الاستخراج بمساعدة الموجات فوق الصوتية للحصول على مستخلصات مائية. تم اختيار خلايا سرطان الجلد (A375) لتقييم المزيد من التأثيرات المضادة للسرطان. تم تعريض خلايا A375 لعمليات تكاثر الخلايا وتشكيل المستعمرات، واستماتة الخلايا، وإيقاف دورة الخلية، وهجرة الخلايا، واختبارات غزو الخلية متبوعة بمسارات جزيئية باستخدام تفاعل سلسلة البلمرة في الوقت الحقيقي (qPCR) وتولد الأوعية في الجسم الحي باستخدام يرقات الزرد (*Danio rerio*). بالإضافة إلى ذلك، تم تقييم جميع ال PBs الأربعة أيضًا لتأثيرها على السمية باستخدام أجنة الزرد. تم تحديد المركبات المضادة للسرطان من PB بشكل افتراضي باستخدام اللوني السائل - مطياف الكتلة (LC-MS) و كروماتوغرافيا السائل - مطياف الكتلة (LC-MS) على أساس نهج التمثيل الغذائي. تم بعد ذلك إرساء جميع المركبات التي تم تحديدها على مركب BCL-2 و cyclin B / CDK1 و VEGF و NM23 البلوري للتنبؤ بتفاعل بروتين ليجاند. وأخيراً، طورت الدراسة أيضًا نموذج انحدار تم التحقق منه باستخدام مطياف تحويل فورييه بالأشعة تحت الحمراء (FT-IR) للتنبؤ بالنشاط المضاد للسرطان لمستخلص PB الجديد لضمان جودته كعامل مضاد للسرطان فعال. أظهرت جميع مستخلصات PB سمية خلوية كبيرة على خلايا A375. كشف التحليل الإضافي أن PB-A و PB-C و PB-D كان له تأثير مثبت جيد على تكاثر الخلايا وتشكيل المستعمرة ودورة الخلية وهجرة الخلايا وغزو الخلايا وتولد الأوعية ومحفز موت الخلايا المبرمج. كشفت PB-A و PB-C و PB-D موت الخلايا المبرمج من خلال المسار الجوهري، مما أدى إلى إيقاف دورة الخلية في المرحلة G2 / M عن طريق خفض تنظيم cyclin B و CDK1. تم إظهار PB-A و PB-C و PB-D على المستوى الجزيئي لقمع NM23 و E-cadherin و MMP2 و MMP9. أظهر تقييم السمية العيوب النمائية الشكلية التي تسببها جميع مقتطفات PB الأربعة مثل قسم الدماغ المشوهة، العمود الفقري الملتوي مع التشوهات في الجسيدات و notochord، التشوهات في الأنسجة الرخوة (الكيس المحي، وذمة التامور والمثانة). بالإضافة إلى ذلك، تبين أن مستخلصات PB تؤثر على أنظمة القلب والأوعية الدموية من خلال وجود وذمة القلب، مما يقلل من معدل ضربات القلب وتدفق الدم. كانت قيم LC50 عند 96 ساعة بعد التخصيب (hpf) أقل من 100 ميكروجرام / مل. اقترب التحليل الكيميائي من مقتطفات PB من خلال استقلاب GC-MS و LC-MS الذي تم تحديده 4-ol-3-4-androsten-17-dione، 7-cholest-3-ol-en، حمض الغاليك، isongifolol، mangiferin وبروبافينون كمبادئ فعالة. تنبأت نتائج الالتحام بتفاعل المبادئ النشطة مع مركب BCL-2 و cyclin B / CDK1 و VEGF و NM23 بشكل رئيسي عن طريق التفاعل مع رهاب الماء مع بقايا البروتين. أظهرت نتائج الإرساء ألفة ربط جيدة للهياكل البلورية لمركب BCL-2 و cyclin B / CDK1 و VEGF و NM23 لـ 4-ol-3-4-androsten-17-dione- و 7-en-cholest-3-ol-en، 4-dimethyl-، 4، 3beta-ol، acetate و mangiferin، مما يشير إلى أدوار هذه المركبات كعوامل مضادة للسرطان محتملة. علاوة على ذلك، أنتجت الدراسة أيضًا نموذجًا إحصائيًا تم التحقق منه للتنبؤ بالنشاط المضاد للسرطان لمستخلصات PB الجديدة. أخيراً، كشفت الدراسة عن نشاط مضاد للسرطان من *Hystrix brachyura bezoar*، والمبادئ النشطة، وآلية العمل في المختبر وتأثير السمية في الجسم الحي باستخدام نموذج الزرد (زبيرا فيش) لأول مرة. تدعم نتائج هذه الدراسة أيضًا الادعاءات التقليدية لاستخدام *Hystrix brachyura bezoar* كعامل مضاد للسرطان في ماليزيا.

APPROVAL PAGE

The thesis of Al'aina Yuhainis Bt Firus Khan has been approved by the following:

Ridhwan Abdul Wahab
Supervisor

Qamar Uddin Ahmed
Co-Supervisor

Alfi Khatib
Co-Supervisor

Norafiza Zainuddin
Internal Examiner

Shaharum Shamsuddin
External Examiner

Nor Fadilah Rajab
External Examiner


DECLARATION

I hereby declare that this thesis is the result of my own investigations, except where otherwise stated. I also declare that it has not been previously or concurrently submitted as a whole for any other degrees at IIUM or other institutions.

Al'aina Yuhainis Bt Firus Khan

Signature

Date



INTERNATIONAL ISLAMIC UNIVERSITY MALAYSIA

**DECLARATION OF COPYRIGHT AND AFFIRMATION OF
FAIR USE OF UNPUBLISHED RESEARCH**

**ANTICANCER STUDY OF PORCUPINE (*Hystrix brachyura*) ON
BEZOAR MELANOMA AND THE IDENTIFICATION OF ITS
ACTIVE COMPOUNDS**

I declare that the copyright holders of this thesis are jointly owned by the student and IIUM.

Copyright © 2020 Al'aina Yuhainis Firus Khan and International Islamic University Malaysia. All rights reserved.

No part of this unpublished research may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise without prior written permission of the copyright holder except as provided below

1. Any material contained in or derived from this unpublished research may be used by others in their writing with due acknowledgement.
2. IIUM or its library will have the right to make and transmit copies (print or electronic) for institutional and academic purposes.
3. The IIUM library will have the right to make, store in a retrieved system and supply copies of this unpublished research if requested by other universities and research libraries.

By signing this form, I acknowledged that I have read and understand the IIUM Intellectual Property Right and Commercialization policy.

Affirmed by Al'aina Yuhainis

.....
Signature

.....
Date

ACKNOWLEDGEMENTS

Bismillahirrahmanirrahim. First and foremost, I thank Allah for giving me the strength and endowing me with health, patience, and knowledge to complete this work. Nothing can be compare to Allah gifts. Almighty ALLAH are the most merciful in providing me the opportunity to have worked with great people who have contributed in various ways to my research and thesis that deserve special mention.

I am thankful to my parents (Firus Khan and Rodziah) for their endless support and love. They are the most wonderful *makhluk* from Allah to me. No words can describe how much I appreciate their loves and supports from birth until now. I wont even have the capability to finish this study without them. I am grateful to my siblings (Farezal, Fazlizul, Fakhru Ikhzan, Fakhru Afdhal, Maslinda, Masliza and Aliyaa. When I am far from home, they give comfort, when am in need of advice, they give beautiful advice, when I need shoulder to cry on, they lend theirs. Thank you as well for taking care of mak and abah when I am far from home.

Next I am thankful to my SV, Assoc.Prof.Dr Ridhwan for accepting me as his students. Am am in deed grateful for all the guidance, advice, about study, life,and the reminders about us here in *dunya* and *akhirah*. Syukran jazilan to my co-SV Dr Qamar Uddin Ahmed, Dr Alfi Khatib, Dr Mardhiah Mohamad and Dr Zalikha Ibrahim for guiding me in my labworks, my articles writing, my data analysis and molecular docking. I am grateful to Allah for giving me the opportunity to meet them. I am lucky enough to meet and having them as my co SV. They really eased my obstacles in my study.

Doing PhD is really tough, but these wonderful *sahabat* make the journey full of joy despite the hurdle. Thank you so much Ayyah, Syamimi, Rosyafirah, Aqilah, Nippun, Alin, Zahirah, Aishah, Ket, Karen for being there with me, listening to my nagging, sharing the difficult times with me, wiping my tears with joys and just for being there for me when I need most. I love you all *lillahitaala* and I pray our friendship remains until *Jannah*. To my labmates, Shakirah, Faizah, Habibah, Hatim, Akilah, Nippun, Salleh, Ahmed and Suga. Not to forget the FYP students whom helped me the most to finish my research faster than usual (Nadzif, Husna, Farahin and Rabbani). I appreciate each one of you for your helps, guidance, advice and supports during my study. I pray Allah ease you matter in your future endeavour.

Special mention to PG KAHS staff Kak Intan, kak Farhana, Kak Hidayah and CPS staff Kak Jihan. Thank you for always reminding me important dates, events, information, giving tips how to deal with postgrad life and always smile when I am facing difficulties. Believe me, nobody (PG KAHS student) can finish their study without all of you and for that I am grateful to Allah.

Thank you as well to few people whom their name shouldn't be mention here. Honestly, I suffered a lot for the difficulties that all of you put me through, however I am thankful for that. I learned how to be strong, bold, determine, more positive and more mature because of you. Nothing happen without Allah permission and I believe, the tough phase is something that I learned most. For that, thank you and please be healthy.

TABLE OF CONTENTS

Abstract	ii
Abstract In Arabic	iii
Approval Page	iv
Declaration	v
Acknowledgements	vii
Table of Contents	viii
List of Tables	xiii
List of Abbreviations	xviii
List of Symbols	xix
CHAPTER ONE: INTRODUCTION	1
1.1 Background of The Study	1
1.2 Problem Statement and Significance of The Study.....	3
1.3 Research Questions	5
1.4 Objectives.....	6
1.1.1 General Objective.....	6
1.1.2 Specific Objectives.....	6
1.5 Research Hypotheses	6
CHAPTER TWO: LITERATURE REVIEW	8
2.1 Natural Product	8
2.1.1 Traditional Medicine	9
2.1.2 Importance of Animal-Based Research.....	13
2.2 Porcupine Bezoar	19
2.2.1 <i>Hystrix Brachyura</i>	19
2.2.2 Medicinal Properties of Bezoar	21
2.2.3 Issues on Porcupine Bezoar.....	22
2.3 Cancer	23
2.3.1 Multi-Step Carcinogenesis	24
2.3.2 Hallmarks of Cancer.....	25
2.3.3 Melanoma.....	27
2.3.4 Melanoma Treatment	28
2.4 Targeted Cellular Process And Molecular Pathway	33
2.4.1 Apoptosis.....	33
2.4.1.1 Apoptosis Cellular Process	33
2.4.1.2 Apoptosis Signalling Pathway	35
2.4.2 Cell Cycle	37
2.4.2.1 Cell Cycle Phases and Checkpoints	38

2.4.2.2 Cell Division Signalling Pathway	39
2.4.3 Angiogenesis	42
2.4.3.1 Pathological of Angiogenesis.....	43
2.4.3.2 VEGF Signalling Pathway	44
2.4.4 Metastasis	46
2.4.4.1 Metastasis Cellular Process.....	46
2.4.4.2 Role of MMP-2, MMP-9, E-Cadherin And NM23.....	48
2.5 Reverse Transcriptase - Quantitative Polymerase Chain Reaction.....	50
2.6 <i>Danio rerio</i> Embryo As Model For Toxicity And Angiogenesis Study	55
2.7 Metabolomics	59
2.8 Molecular Docking.....	63

CHAPTER THREE: ANTICANCER ACTIVITY OF DIFFERENT PORCUPINE BEZOAR EXTRACT ON MELANOMA CELL (A375) 66

3.1 Introduction	66
3.2 Materials.....	67
3.3 Methodology	67
3.3.1 PB Extract Preparation	67
3.3.2 <i>In Vitro</i> Cytotoxicity Analysis	68
3.3.2.1 Cell Culture Technique	68
3.3.2.2 Measuring Cells Viability	69
3.3.2.3 Cells Seeding For Assay	69
3.3.2.4 Cells Optimization	70
3.3.2.5 Cytotoxicity Assay of on Cancer Cells	70
3.3.2.6 Cytotoxicity on Normal HDF.....	71
3.3.3 Growth Inhibition Assay	71
3.3.4 Annexin-V Assay	72
3.3.5 Cell Cycle Assay	72
3.3.6 Migration Assay	73
3.3.7 Invasion Assay	73
3.3.8 Colony Formation Assay.....	74
3.3.9 Gene Expression Analysis Using Reverse Transcriptase Quantitative Polymerase Chain Reaction (RT- qPCR).....	75
3.3.9.1 RNA Extraction.....	75
3.3.9.2 Measuring RNA Purity and Concentration	75
3.3.9.3 RNA Integrity Assay.....	76
3.3.9.4 cDNA Transcription.....	76
3.3.9.5 Primer Selection	77
3.3.9.6 Optimization of Annealing Temperature and Melt Curve For Primer	79
3.3.9.7 Optimization of Primer Efficiency with The Standard Curve.....	80
3.3.9.8 Evaluation of Targeted Genes Using RT-qPCR	81
3.3.10 Angiogenesis Assay	82
3.3.10.1 Maintenance of <i>Danio Rerio</i>	82
3.3.10.2 Spawning of <i>Danio Rerio</i> and Embryo Care	82
3.3.10.3 Whole-Mount Alkaline Phosphatase Staining	82

3.4 Result.....	84
3.4.1 Extraction Yield	84
3.4.2 Cytotoxicity of Aqueous PBs Extracts on A Panel of Cells Line	84
3.4.3 Cytotoxicity on Normal Human Dermal Fibroblast (NHDF) ...	86
3.4.4 Effects of PBs Extract on Cell Growth Activity of A375 Cells	87
3.4.5 Effects of PBs Extracts on Cell Cycle Activity of A375 Cells .	90
3.4.6 The Effects of PBs Aqueous Extracts on The Apoptotic Activity of A375 Cells.	93
3.4.7 Effects of PB Extracts on Anti-Metastatic Activity on A375 ...	96
3.4.8 Effects of PB Extracts of Antiangiogenesis on Embryo <i>Danio</i> <i>rerio</i>	102
3.5 Discussion	104

CHAPTER FOUR: EMBRYO *Danio rerio* TOXICITY OF DIFFERENT PORCUPINE BEZOARS EXTRACTS..... 112

4.1 Introduction	112
4.2 Methodology	113
4.2.1 Extract Preparation And Treatment Procedure	113
4.2.2 Teratogenicity Index (TI) Evaluation.....	114
4.2.3 Evaluation of Lethal And Sublethal Toxicity	114
4.3 Result.....	115
4.3.1 Median Lethal Concentration, Median Effective Concentration and Teratogenicity Index	115
4.3.2 Effect of PBs Aqueous Extracts on Spinal And Body Shape....	117
4.3.3 Effect of PBs Aqueous Extracts on Somite And Notochord....	117
4.3.4 Effect of PBs Aqueous Extracts on Developmental Defects In <i>Danio rerio</i> Embryos Brain	120
4.3.5 Effects of PBs Aqueous Extracts on Soft Tissues (Yolk Sac, Pericardial Oedema And Swim Bladder).....	122
4.3.6 Effect of PBs Aqueous Extracts on Cardiovascular Function...	124
4.4 Discussion	126

CHAPTER FIVE: PROFILING OF PORCUPINE BEZOAR USING METABOLOMICS APPROACH 130

5.1 Introduction	130
5.2 Methodology	131
5.2.1 Porcupine Bezoar Extract Preparation For Metabolomics	131
5.2.2 Cytotoxicity of Porcupine Bezoar Extracts On A375 Cells	131
5.2.3 Metabolites Profiling Using Gas Chromatography - Mass Spectrometer (GCMS)	132
5.2.4 Derivatization of Sample.....	132
5.2.5 Analytical Method and Data Processing	132
5.2.6 Metabolites Profiling Using Q-TOF Liquid Chromatography-Mass Spectrometry (Q-TOF LC-MS)	133
5.3 Result.....	134
5.3.1 Extraction Yield	134

5.3.2 Melanoma Cytotoxicity Evaluation	134
5.3.3 Metabolites Profiling Using GC-MS.....	136
5.3.4 Multivariate Data Analysis.....	140
5.3.5 Metabolites Profiling Using Q-TOF LC-MS	144
5.3.5.1 Multivariate Data Analysis of Q-TOF LC-MS Data <i>Hystrix</i> <i>Brachyura</i> Bezoar	144
5.3.6 Structure Elucidation Using Fragmentation Pathway	147
5.4 Discussion	152

CHAPTER SIX: IN SILICO EVALUATION OF IDENTIFIED ACTIVE COMPOUNDS WITH TARGET PROTEINS

6.1 Introduction	158
6.2 Methodology	159
6.2.1 Ligand Preparation	159
6.2.2 Protein Preparation	159
6.2.3 Docking And Analysis with Ligplot	160
6.3 Result.....	162
6.3.1 Molecular Interaction of Target Compounds with Bcl-2 Protein	162
6.3.2 Molecular Interaction of Target Ligands with Cyclin B/CDK1 Complex Protein	169
6.3.3 Molecular Interaction of Target Ligands with VEGF Protein ..	175
6.3.4 Molecular Interaction of Target Ligands with NM-23 Protein .	181
6.4 Discussion	186

CHAPTER SEVEN: CHARACTERIZING PORCUPINE BEZOAR FUNCTIONAL GROUPS USING METABOLOMICS APPROACH..... 200

7.1 Introduction	200
7.2 Methodology	201
7.2.1 Analytical Method.....	201
7.2.2 Data Processing and Multivariate Data Analysis	201
7.2.3 Validation and Predictive Capability of Multivariate Data Analysis.....	202
7.2.4 Statistical data analysis.....	202
7.3 Result.....	202
7.3.1 Multivariate Data Analysis.....	204
7.3.2 Validation And Predictive Capability of MVDA.....	207
7.4 Discussion	208

CHAPTER EIGHT: CONCLUSION AND FUTURE RESEARCH RECOMMENDATION 212

REFERENCE 216

APPENDIX A: LIST OF MATERIALS USED IN THIS STUDY..... 243

APPENDIX B: APPROVAL FROM PERHILITAN 250

APPENDIX C: GROWTH CURVE OF CELLS USED IN THIS STUDY... 252

APPENDIX D: GENE EXPRESSION ANALYSIS USING REVERSE TRANSCRIPTASE QUANTITATIVE POLYMERASE CHAIN REACTION (RT-QPCR).....	253
APPENDIX E: LCMS SPECTRA OF ALL EXTRACTS	268
APPENDIX F: LC-MS FRAGMENTATION SPECTRA	269
APPENDIX G: FRAGMENTATION PATHWAY OF ANTICANCER COMPOUNDS IDENTIFIED USING Q-TOF-LCMS	271
APPENDIX H: MOLECULAR DOCKING OF 5-FU WITH THYMIDYLATE SYNTHASE PROTEIN.....	285
APPENDIX I: LIST OF CONFERENCES AND PUBLICATION	286



LIST OF TABLES

<u>Table No</u>		<u>Pages</u>
2.1	List of traditional medicine system	10
2.2	List of animal and purposed used in traditional medicine.	14
3.1	Primer used in this study	78
3.2	Reaction components for PCR reaction in each well.	80
3.3	Protocol for standard cycling.	80
3.4	Extraction yields of all PB using water as solvents.	84
3.5	Median inhibition concentration of PBs aqueous extracts on panel of cancer cells line with UT and 5-FU as controls	85
4.1	LC ₅₀ , EC ₅₀ and TI values as derived from the concentrations-response curves for all PBs extract.	116
5.1	Extraction yield of PB according to solvent.	136
5.2	Effect of solvent median concentration of cytotoxic assay on A375 cells of PB extract.	136
5.3	Tentative compounds identified in the PB extracts through GC-MS analysis	139
5.4	Tentative anticancer compounds identified in the PB methanol extract through Q-TOF LC-MS/MS fragmentation using positive ionisation	149
6.1	Protein used in molecular docking analysis.	161
6.2	Molecular interaction results of Bcl-2 protein with the known anticancer drugs (5-FU) and the active ligands.	163
6.3	Molecular interaction results of cyclin B/CDK1 complex protein with the known anticancer drugs (5-FU) and the active ligands identified using GC-MS	170
6.4	Molecular interaction results of VEGF protein with the known anticancer drugs (5-FU) and the active ligands identified using GC-MS	176
6.5	Molecular interaction results of NM 23-H1 protein with the known anticancer drugs (5-FU) and the active ligands identified.	182
7.1	Infrared spectral assignments of PB extracts.	204
7.2	Measured and predicted cytotoxic activity of 100% methanol PB extract obtained from unknown sources in Peninsular Malaysia.	208

LIST OF FIGURES

<u>Figure No</u>		<u>Pages</u>
1.1	Flowchart of this study	7
2.1	<i>Hystrix brachyura</i> or known as Malayan porcupine	20
2.2	Pendant of porcupine bezoar.	22
2.3	Cancer development process	24
2.4	The Hallmarks of cancer.	27
2.5	Morphology schematic of cells that undergo apoptosis	34
2.6	The schematic diagram representing the main molecular pathways of apoptosis, the extrinsic and intrinsic pathways.	37
2.7	Phase of cell cycle with its regulatory Cyclin/CDKs complex	42
2.8	Comparison early process of angiogenesis between sprouting and intussusceptive angiogenesis	44
2.9	Schematic diagram showing the different stages of cancer metastasis.	47
2.10	DNA replication in cells	51
2.11	Amplification curve, melt curve and standard curve of RT-qPCR.	54
2.12	Lateral view of morphological end points for toxicity assessment at 120 hpf	57
2.13	Alkaline phosphatase staining at 72 hpf revealing the vasculature of <i>Danio rerio</i>	58
2.14	Flowchart of the metabolomics study.	60
2.15	Molecular docking interaction between ligand and receptor	65
3.1	Image of porcupine bezoar used in this study	68
3.2	Log dose response of all PBs aqueous extract against cancer cells.	85
3.3	The effects of PBs extracts on NHDF cells growth	87
3.4	The effects of PBs extracts on A375 cells growth.	89
3.5	The effects of PBs extracts on A375 cell cycle arrest at 72 h incubation	91
3.6	Effects of PBs extracts on the cell cycle arrest gene mRNA expression of A375.	92
3.7	The effects of PBs extracts on apoptosis analysis.	94

3.8	Effects of PBs extracts on the targeted apoptosis gene mRNA expression of A375.	95
3.9	Anti-migration effect of PBs extracts against A375 cells.	97
3.10	Anti-invasion effect of PBs extracts against A375 cells.	99
3.11	The effects of PBs extracts on A375 colonies formation	100
3.12	Effects of PBs extracts on the targeted metastasis gene mRNA expression of A375	101
3.13	The effects of PBs extracts on <i>Danio rerio</i> embryo angiogenesis assay	103
3.14	Porcupine bezoar anticancer activity mechanism of action in A375 cells in this study.	111
4.1	Log concentration-response curves for malformation and mortality of <i>Danio rerio</i> embryos and larvae at different hpf in different PBs extract concentrations	116
4.2	The effect of 1.25 mg/mL of PBs aqueous extract treated on embryos/larvae spine and body shape morphology at 96 hpf.	118
4.3	Effect of PBs on somites and notochord. The images represented the morphology of embryos at 96 hpf with PC and NC as control	119
4.4	The PBs extract effect on gross morphology of brain section and craniofacial.	121
4.5	Morphology of embryo <i>Danio rerio</i> soft tissues upon treatment with PBs extract.	123
4.6	Effects of cardiovascular function upon PBs extract exposure. The observation was recorded at 72 hpf.	125
5.1	GC-MS chromatogram of metabolites in 0% (A), 25% (B), 50% (C), 75% (D) and 100% (E) of methanol: water dilution of PB extract	138
5.2	The score plot of PB extracts (A, B, C, D and E) based on the OPLS obtained using GC-MS.	141
5.3	The summary of fit of OPLS models of the PB extracts (A, B, C, D and E) analysed using the GC-MS.	141
5.4	Permutation of OPLS model of PB extracts (A, B, C, D and E) analyzed using GC-MS.	143
5.5	The loading scatter plot OPLS models of PB methanol extract using GC-MS.	143
5.6	Metabolites identified as potential active anticancer agents from PB methanol extract.	144
5.7	The score plot of different extracts of PB extracts (A, B, C, D and E) based on the OPLS obtained using Q-TOF LC-MS positive ionization.	146

5.8	The summary of fit of OPLS models of the PB extracts analysed using Q-TOF LC-MS positive ionization.	146
5.9	The loading scatter plot of OPLS models of the PB methanol extract analysed using Q-TOF LC-MS with positive ionization	147
5.10	Metabolites identified as potential active anticancer agents from PB methanol extract.	148
6.1	The 2D diagram of the protein-ligand interactions with the residues of Bcl-2 and ligands	165
6.2	The 2D diagram of the protein-ligand interactions with the residues of the Bcl-2 with GC-MS identified ligands	166
6.3	The 2D diagram of the protein-ligand interactions with the residues of Bcl-2 and Q-TOF LC-MS identified ligands	168
6.4	The 2D diagram of the protein-ligand interactions with the residues of cyclin B/CDK1 complex and ligands	171
6.5	The 2D diagram of the protein-ligand interactions with the residues of the cyclin B/CDK1 complex with GC-MS identified ligands	172
6.6	The 2D diagram of the protein-ligand interactions with the residues of cyclin B/CDK1 complex and LC-MC identified ligands.	174
6.7	The 2D diagram of the protein-ligand interactions with the residues of VEGF and ligands.	177
6.8	The 2D diagram of the protein-ligand interactions with the residues of the VEGF with GC-MS identified ligands.	179
6.9	The 2D diagram of the protein-ligand interactions with the residues of VEGF and LC-MC identified ligands	180
6.10	The 2D diagram of the protein-ligand interactions with the residues of NM23-H1 and ligands.	183
6.11	The 2D diagram of the protein-ligand interactions with the residues of the NM23-H1 with GC-MS identified ligands	185
6.12	The 2D diagram of the protein-ligand interactions with the residues of NM23-H1 and LC-MC identified ligands.	186
6.13	The superimposed 3D diagram showing the binding site of ligands on Bcl-2 protein.	190
6.14	The superimposed 3D diagram showing the binding site of ligands on cyclin B/CDK1 complex	193
6.15	The superimposed 3D diagram showing the binding site of ligands on VEGF protein	195

6.16	The superimposed 3D diagram showing the binding site of ligands on NM23-H1 protein	198
7.1	FT-IR spectra of porcupine bezoar extracted using different dilution of solvents.	203
7.2	The FT-IR score scatter plot porcupine bezoar extracted using different dilution of solvents based on the partial least squares (OPLS).	205
7.3	Line loading plot of the PB methanol extract FT-IR orthogonal partial least squares model	206
7.4	Permutation OPLS model of PB extracts (A, B, C, D and E) analyzed using FT-IR.	207
7.5	Prediction versus observation IC_{50} values from all PB extracts.	207



LIST OF ABBREVIATIONS

ATCC	American type culture collection
A375	Melanoma cancer cells
BLAST	Basic local alignment search tool
Bax	Bcl-2-associated X protein
Bcl2	B-cell lymphoma 2
BID	BH3-interacting-domain death
CASPASE	Cysteine aspartic acid protease
CDKs	Cyclin-dependent kinases
CDKI	Cyclin-dependent kinases inhibitors
CDNA	Complementary DNA
CGM	Complete growth media
CIP/KIP	CDK Interacting protein/kinase inhibitor protein
CO ²	Carbon dioxide
dIH ₂ O	Dionize distilled water
DNA	Deoxyribonucleic acid
DMEM	Dulbecco's Modified Eagle Medium
DMSO	Dimethyl sulphoxide
E	Efficiency
EC ₅₀	Median efficient concentration
FADD	Fas-associated death domain protein
FasL	Fas ligand
FBS	Fetal bovine serum
IC ₅₀	Median inhibition concentration
LC ₅₀	Median lethal concentration
GAPDH	Glyceraldehyde-3-phosphate dehydrogenase
ACTB	Beta actin
GC	Guanine-cytosine (DNA base pairing)
LC50	Lethal concentration
mRNA	messenger RNAs
MIQE	Minimum information for quantitative polymerase chain reaction publication experiments
NCBI	National centre for biotechnology
NC	Negative control
NHDF	Normal human dermal fibroblast
PB	Porcupine bezoar
PBS	Phosphate buffer saline
PCR	Polymerase chain reaction
PC	Positive control
PI	Propidium Iodide
qPCR	Quantitative polymerase chain reaction
RNA	Ribonucleic acid
RT	Real-Time
UT	Untreated
7AAD	7-Aminoactinomycin D

LIST OF SYMBOLS

α	Alpha
β	Beta
Δ	Delta
Cq	Quantification cycle
Ct	Threshold cycle
g	Gram
G	Gravity
G	Gap
hpf	Hours post-fertilization
M	Mitosis
S	Synthesis
μ l	Microliter
μ g/ml	Microgram per millilitre
$^{\circ}$ C	Degree Celsius
%	Percent
-	To
>	More than
<	Less than
\pm	Plus-minus
x	Times
=	Equal to
*	Statistical significance denotation

CHAPTER ONE

INTRODUCTION

1.1 BACKGROUND OF THE STUDY

Natural products have played an important aspect in life since ancient time as it was integrated as natural resources into daily life for food and remedies for various reasons. Natural product-based medicines have been widely used and practised in Ayurveda, Traditional Chinese Medicine, Traditional Malay Medicine, Greek and Roman Medicine systems, and many more (Cragg et al., 2009; Ji et al., 2009; Rocha et al., 2001). Currently, researchers are exploring scientific findings of utilisations of natural products resources as alternative medicines to treat various illnesses, including cancer (Lahlou, 2013). Natural product had been the single most productive source of leads for the development of drugs where more than 100 new anticancer drugs were in clinical development (Harvey, 2017).

Besides, apart from using herbal or plant for remedies, traditional medicine had been using animal-based remedies for various medical condition. The phenomenon of using animal-based therapy exist until today due to deep rooted-historical origins and species geographical distribution (Alves & Rosa, 2005). The usage of animal-based includes body parts for instance fat, bile, brain, bones, fin, claws, horn, feather, scales, teeth, skin, eggs, milk, blood, bezoar or whole animal (Costa-neto, 2005; Alves et al., 2011; Ma et al., 2015). Despite the practice of animal-based worldwide, researchers had been neglecting the studies of the therapeutic use of animals and animal parts compared to plants. It is crucial to investigate the practice to verify the traditional claims, study the medicinal efficacy and its toxicity. Since there is very limited information on

animal-based medicine, the public tends to believe the history and traditional belief instead of using modern medicine.

The present study will investigate the usage of the animal derived-natural product, namely porcupine bezoar (PB) as alternative medicine/ supplement to treat cancer. Porcupine bezoars are concretions of undigested organic or inorganic compounds found in the gastrointestinal tract of porcupine (Mori & Sforzi, 2013). Bezoars are categorised into four classes which include phytobezoars, trichobezoar, lactobezoar and pharmacobezoar (Sanders, 2004). The bezoar is phytobezoar which is commonly composed of undigested fibres, skins, seeds, leaves, roots or stems of the plant depending on what the porcupine consumed (El Fortia et al., 2006; Sanders, 2004). Bezoar once was believed to be universal antidotes or panacea that can cure all kinds of deadly diseases or illness (Duffin, 2013; Barroso, 2014).

In Malaysia, the most common porcupine is *Hystrix brachyura*, a species of rodent from the Hystricidae family and is well known as a Malayan porcupine or Himalayan porcupine (Magintan et al., 2017). The PB is known to local aborigines as “*Geliga Landak*” and considered quite famous compared to other animal bezoar. The PB is considered as exotic medicine as porcupine numbers are decreasing and under the protection of Malaysian law. Moreover, PB is a rare material as less number of porcupine forms the bezoar. Therefore, poachers and traders tend to sell it implausibly at higher prices. Since the 8th century, numerous claims and myth were lingering around bezoar claiming it as the prince of antidotes capable of curing any illness. According to Shan et al., (2019) in Malaysia, the most famous story regarding PB comes from Chinese folklore, where they believed PB capable of treating cancer and as supplements post-cancer therapy. Despite the traditional claims, the anticancer properties of PB have never been investigated thoroughly. Therefore present study investigates PB

metabolomics profile, anticancer properties and its mechanism of action on melanoma, a malignant skin-derived cancer.

Melanoma arises from melanocytes, specialised pigmented cells that are found predominantly in the skin (Gray-schopfer, Wellbrock, & Marais, 2015). Melanocytes are specialised pigmented cells that are found predominantly in the skin and eyes, where they produce melanin, the pigments responsible for skin and hair colour (Tas, 2012). Melanoma has the greatest metastatic potential among other skin cancer. Though melanoma is diagnosed around 4% compared to all skin cancer, it contributed to 75% of skin cancer mortality (Arrangoiz et al., 2016). Moreover, metastatic melanoma has a poor five-year survival prognosis with increasing incidence and mortality rate keep increasing (Rigel & Carucci, 2000; Chakraborty et al., 2013). Melanoma is an extremely aggressive type of cancer with high metastatic potential and notoriously high resistance to cytotoxic agents probably due to melanocytes originating from highly motile cells that have enhanced survival properties (Gray-schopfer et al., 2015). Furthermore, the study reported for the past 40 years, limited progress has been made in metastatic melanoma treatment (Chakraborty et al., 2013).

1.2 PROBLEM STATEMENT AND SIGNIFICANCE OF THE STUDY

Traditional medicine is the oldest health care used since ancient time to manage prevention, treating and curing disease or any medical condition. Each part of the world has its deep-rooted origins of traditional medicine to combat various health and life-threatening medical condition (Still, 2003). The traditional medicine advances have developed over the past centuries concerning methods of preparation, extraction, herbs selection, medicinal materials identification, the best time for obtaining different natural resources and prescription (Yuan et al., 2016). Though modern medicine had been well

developed, there are many societies still used traditional medicine as complementary or as alternative medicine.

In recent decades, the interest in evaluating the therapeutic effects of natural products has been increasing. However, most studies focused mainly on herbal and plants based medicine. Since traditional medicine includes the usage of materials from plants and animal, there is a huge gap in understanding traditional medicine in the past. Opposite to herbal medicine, animal-based medicine had less information. The preparation and prescription depend on traditional practice and belief rather than scientifically studied (Still, 2003). This situation caused excessive hunting by poachers as most of the animal-based medicine used exotic and wild animal hence leading to rapid extermination of numerous animal species.

According to Molur et al., (2005), *Hystrix brachyura* species in 2005 was enlisted as endangered animal in Bangladesh, and near threatened in India and Nepal. The same report mentions that the porcupine population had been declining by more than 10% for the last 20 years. The main reason for population declining are habitat loss and hunting for medicinal purpose. The porcupine population was declining in Malaysia as well it is now under the Malaysian government protection of Wildlife Act (1972). Although porcupines are under Malaysian law protection, however, the aborigines were permitted to continue hunting porcupine as it is a tradition passed down from their ancestor to consume and used the porcupine for medicinal purposes (Shan et al., 2019). Most of the PB was obtained from the aborigines and supplied to local traditional Chinese medicine store. The selling price of the PB in the Malaysian market is around RM 300-RM 1000 for 500mg of PB, due to difficulty in obtaining it. This circumstance

creates an overflow of fake PB and increasing numbers of porcupine killed for easy money.

There is an urgency to verify traditional claims of PB anticancer properties, as there is no scientific information on PB medicinal benefits to support. The present study will investigate PB anticancer properties from different porcupines and elucidate its mechanism of action on melanoma cells (A375) using Reverse Transcription Quantitative Polymerase Chain reaction (RT-qPCR). Toxicity profile of different PBs will be conducted using *Danio rerio* embryo to evaluate its toxicity profile. The PB metabolomics profile using gas chromatography-mass spectrometry (GC-MS) and quadrupole time of flight liquid chromatography-mass spectrometry (QTOF LC-MS) will be evaluated as well to determine possible active compounds and characterize PB extracts. The potential active compounds interaction with targeted proteins will be analysed using molecular docking approach. A validated regression model using FT-IR based fingerprinting will be developed to predict PB anticancer activity on melanoma cells (A375).

1.3 RESEARCH QUESTIONS

The present study aims to answer the following questions:

- Do PBs aqueous extract safe for consumption?
- Do different PBs aqueous extracts have anticancer properties on melanoma?
- What are the active compounds contributing to anticancer activities?
- Do active compounds identified in this study interact with similar targeted proteins?
- Do PB extracts from different forest have a similar anti-cancer effect?