

*IN SILICO* STUDY, CLONING AND FUNCTIONAL  
ANALYSIS OF CjS8 PROTEIN FROM  
*Campylobacter jejuni*

BY

HATIM ABDULLAH YOUSEF NATTO

A thesis submitted in fulfilment of the requirement for the  
degree of Doctor of Philosophy

Kulliyyah of Allied Health Sciences  
International Islamic University Malaysia

JULY 2020

## ABSTRACT

In Gram-negative bacteria, protein secretion plays an important part in pathogenesis. Secretory proteins perform a variety of important role for bacterial survival in the environment and its involvement to cause disease to human. We have identified specific surface protein of *Campylobacter jejuni*. Genomic and protein analysis using bioinformatics tools on that protein reveals the presence of signal peptide at N-terminal of its peptide sequence, thus signifies this protein is highly potential acted as secreted protein. The size of the protein was calculated as 24.1 kDa. The availability of complete genome sequences of *C. jejuni* has allowed this study to make predictions about the composition of bacterial secreted protein that has good similarity in their sequence homology. The prediction of others *C. jejuni* secreted proteins were performed based on 24.1 kDa and several enterobacteriaceae pathogenic bacteria proteins sequences using a set of internet-based programs, including BLAST, ORF Finder and SignalP v 5. *In silico* analysis in this study identified the secreted protein of S8 family serine peptidase in *C. jejuni* genome and designated as CjS8. To investigate the involvement of CjS8 in the pathogenesis of *C. jejuni* infection, a C-terminal fragment of CjS8 was successful amplified, cloned and expressed using a TOPO expression vector. Rabbit polyclonal serum was raised against the purified recombinant CjS8 protein. The CjS8 null-mutant was constructed in *C. jejuni* by natural transformation and allelic exchange. PCR analysis and immunoblot of whole cell lysates with Ab\_CjS8 (polyclonal antibody against CjS8) showed that CjS8 is naturally expressed in *C. jejuni* but not in the null mutant. In a strain survey on clinical isolates of *C. jejuni*, using the PCR and immunoblot analysis. Data showed that the CjS8 was presence in twenty out of twenty-three clinical isolates. Importantly, this study revealed the functional analysis results, that showed the CjS8 mutant was shown to affect the *C. jejuni* ability to adhere to the host cells (Caco-2 cells). Invasion was also affected by CjS8 mutant strain as well as the biofilm formation and motility. Thus, as a conclusion the CjS8 was the one, among a few secreted proteins described in *C. jejuni* and may represent a novel virulence factor. These results will be important in furthering our understanding of Campylobacter biology and pathogenesis.

## خلاصة البحث

في البكتيريا سالبة الجرام ، يلعب إفراز البروتين دورًا مهمًا في التسبب في المرض. تؤدي البروتينات الإفرازية مجموعة متنوعة من الأدوار المهمة لبقاء البكتيريا في البيئة ومشاركتها في إحداث المرض للإنسان. لقد قمنا بتعريف بروتينًا سطحيًا محددًا للكامبيلوباكتري جيجوني. ان تحليل الجينوم والبروتين باستخدام أدوات المعلوماتية الحيوية على هذا البروتين يكشف عن وجود سيجنال بيتيد في الطرف ان من تسلسل الببتيد، وبالتالي يشير إلى أن هذا البروتين محتمل للغاية ان يعمل كبروتين مُفرز. لقد تم حساب حجم البروتين على أنه 24.1 كيلو دالتون. ان توفر التسلسل الجيني الكامل للكامبيلوباكتري جيجوني اتاح لهذه الدراسة اجراء تنبؤات حول تكوين البروتينات البكتيرية المفزة التي لها تشابه جيد في التجانس الجيني. تم اجراء التنبؤ بالبروتينات الاخرى غير الكامبيلوباكتري جيجوني التي يتم افرازها والتي لها تشابه جيد في التسلسل الجيني. تم اجراء التنبؤ بالبروتينات الأخرى التي يتم افرازها من الكامبيلوباكتري جيجوني استناداً الى 24.1 كيلودالتون والعديد من متواليات البكتريا المعوية المسببة للأمراض وذلك باستخدام مجموعة من البرامج المستندة الى الانترنت، بما في ذلك بلاست و اوين ريدنيج فريم وسيجنال بي. ان التحليل باستخدام السيلكو في هذه الدراسة حدد بروتين مفرز من عائلة اس 8 سيرين بيبتيديز في جينوم الكامبيلوباكتري جيجوني تم تسميته سي جيجوني اس 8. للكشف عن دور سي جيجوني اس 8 في التسبب بالمرض والاصابة بالعدوى بالكامبيلوباكتري جيجوني، تم بنجاح تكوين السي تيرمنل وزراعته و انتاج البروتين باستخدام توبو أكسبريشن فيكتور. تم انتاج مصل ارنب متعدد النسيلة ضد البروتين المنقى من سي جيجوني اس 8. تم انشاء الطفرة الجينية في السي جيجوني اس 8 بواسطة الانتقال الطبيعي والتبادل الأليلي. نتأج تفاعلات سلسلة البوليميز والغشاء المناعي الناتج بتحليل كامل الخلايا مع الاجسام المضادة الناتجة ضد سي جيجوني اس 8، اظهرت أن سي جيجوني اس 8 ينتج طبيعياً في الكامبيلوباكتري جيجوني ولكن لاينتج في الطفرة الجينية فيها. في الدراسة الاستقصائية للسلاطات على عينات طبية للكامبيلوباكتري باستخدام تفاعلات سلسلة البوليميز وتحليل الغشاء المناعي الناتج بتحليل كامل الخلايا، اظهرت البيانات ان سي جيجوني اس 8 موجود في عشرون عزلة من اصل ثلاثة وعشرين عزلة طبية. الأهم من ذلك كشفت هذه الدراسة عن نتأج التحليل الوظيفي حيث اوضحت أن هناك تأثير للسي جيجوني اس 8 الطفرة الوراثية على قدرة الكامبيلوباكتري جيجوني على الالتصاق بخلايا العائل المضيف. كذلك الغزو تأثر بالطفرة الوراثية للسي جيجوني اس 8 وكذلك الحال بالنسبة لكل من تشكيل الأغشية الحيوية والحركة. وهكذا فانه كاستنتاج فان السي جيجوني اس 8 يعتبر واحداً من بين عدد من البروتينات المفزة الموصوفة في الكامبيلوباكتري جيجوني والذي يشكل عامل ضراوة جديد. سوف تكون هذه النتأج مهمة في تعزيز فهمنا لبيولوجيا الكامبيلوباكتري وطرق احداث المرض.

## **APPROVAL PAGE**

The thesis of Hatim Abdullah Yousef Natto has been approved by the following:

---

Ridhwan Abdul Wahab  
Supervisor

---

Mohd Zaki Salleh  
Co-Supervisor

---

Robaiza Zakaria  
Co-Supervisor

---

Mohammad Syaiful Bahari Abdull Rasad  
Internal Examiner

---

Tang Thean Hock  
External Examiner

---

Mohd Nasir Mohd Desa  
External Examiner

---


Jesni Shamsul Shaari  
Chairman

## DECLARATION


I hereby declare that this dissertation is the result of my own investigations, except where otherwise stated. I also declare that it has not been previously or concurrently submitted as a whole for any other degrees at IIUM or other institutions.

Hatim Abdullah Yousef Natto

Signature .....



Date .....



**INTERNATIONAL ISLAMIC UNIVERSITY MALAYSIA**

**DECLARATION OF COPYRIGHT AND AFFIRMATION OF  
FAIR USE OF UNPUBLISHED RESEARCH**

***IN SILICO* STUDY, CLONING AND FUNCTIONAL ANALYSIS  
OF Cjs8 PROTEIN FROM *Campylobacter jejuni***

I declare that the copyright holders of this thesis are jointly owned by the student and IIUM.

Copyright © 2020 Hatim Abdullah Yousef Natto and International Islamic University Malaysia. All rights reserved.

No part of this unpublished research may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise without prior written permission of the copyright holder except as provided below

1. Any material contained in or derived from this unpublished research may be used by others in their writing with due acknowledgement.
2. IIUM or its library will have the right to make and transmit copies (print or electronic) for institutional and academic purposes.
3. The IIUM library will have the right to make, store in a retrieved system and supply copies of this unpublished research if requested by other universities and research libraries.

By signing this form, I acknowledged that I have read and understand the IIUM Intellectual Property Right and Commercialization policy.

Affirmed by Hatim Abdullah Yousef Natto



.....  
Signature

.....  
Date

## DEDICATION

It is my utmost pleasure to dedicate this thesis to the beloved parents, may God have mercy on them, who moved to the mercy of God during the final stage of writing on this thesis. Also this thesis is dedicated to my family, who have been my backbone and words cannot express my deepest gratitude and love. Thank you for your support and patience.



## ACKNOWLEDGEMENTS

First of all, I would like to thank my God Almighty who helped and supported me step by step for me to go through on this study. I wish to express my appreciation and thanks to those who provided their time, effort and invaluable support for this study. A special thanks to my supervisor Associate Professor Dr. Ridhwan Abdul Wahab for his continuous and invaluable support, encouragement, friendship and for that, I will be forever grateful. To my Co-Supervisors, Professor Dato' Dr. Mohd Zaki Salleh (Director, Integrative Pharmacogenomics Institute (iPROMISE), UiTM) and Dr. Robaiza Zakaria (Department of Medical Microbiology and Parasitology, Medical School, USM) for their valuable encouragement and advice in this study. And also not to forget Professor Dato' Dr. Ridzwan Hashim, Dr Ariffin and many more academic staffs in IIUM that helping me directly or indirectly for the successful of this study. One small smiles from your clean heart, means something to me and motivate me to move on.

I would like to express my thanks to all in Biomedical Science Department in particular, Assistant Professor Dr. Hanani Binti Ahmad Yusof @ Hanafi (Head of Department), KAHS postgraduate office and the Dean, Professor Dr. Suzanah Binti Abdul Rahman for their support during my candidature as a Ph.D student in IIUM.

I would like to extend my warm and sincere thanks to my research team-mates, Sister Alaina, Tara, Faizah, Habibah and Shakirah for their friendship, helps, supports, and encouragement.

I wish to express my sincerest appreciation to Saudi Government in particular, Umm Al-Qura University and the Saudi Embassy and Saudi Cultural Attache staffs for their support and providing me this opportunity to complete the Ph.D degree at IIUM.

## TABLE OF CONTENTS

Abstract.....	ii
Abstract in Arabic.....	iii
APPROVAL PAGE .....	iv
DECLARATION .....	v
Copight Page	vi
INTERNATIONAL ISLAMIC UNIVERSITY MALAYSIA.....	vi
DEDICATION .....	vii
ACKNOWLEDGEMENTS .....	viii
Table of contents .....	ix
List of Tables .....	xiv
List of Figures.....	xv
ABBREVIATIONS .....	xix
CHAPTER ONE .....	1
INTRODUCTION .....	1
1.1 BACKGROUND OF THE STUDY.....	1
1.2 PROBLEM STATEMENT AND SIGNIFICANCE OF THE STUDY .	3
1.3 RESEARCH QUESTIONS .....	4
1.4 OBJECTIVES .....	5
1.5 RESEARCH HYPOTHESIS.....	6
CHAPTER TWO .....	8
LITERATURE REVIEW .....	8
2.1 THE GENUS <i>CAMPYLOBACTER</i> .....	8
2.2 CLASSIFICATION OF <i>CAMPYLOBACTER</i> .....	8
2.3 GENOMIC OVERVIEW OF <i>CAMPYLOBACTER</i> .....	9
2.4 ISOLATION AND CULTURE OF <i>CAMPYLOBACTER</i> .....	12
2.5 IDENTIFICATION OF <i>CAMPYLOBACTER</i> .....	13
2.6 DETECTION BY POLYMERASE CHAIN REACTION (PCR) .....	14
2.7 EPIDEMIOLOGY OF <i>CAMPYLOBACTER</i> .....	15
2.8 PATHOGENESIS AND VIRULENCE FACTORS OF <i>CAMPYLOBACTER</i> .....	17
2.8.1 Toxins .....	18
2.8.2 Adhesion.....	18
2.8.3 Invasion .....	19
2.8.4 Intracellular survival.....	21
2.8.5 Trans-epithelial translocation .....	21
2.8.6 Flagella and Motility .....	22
2.8.7 Lipooligosaccharide and capsular polysaccharide .....	23
2.8.8 Biofilm formation.....	24
2.9 GRAM NEGATIVE BACTERIAL PROTEIN SECRETION MECHANISMS .....	25
2.9.1 Type I secretion pathway .....	26
2.9.2 Type II secretion pathway .....	27
2.9.3 Type III secretion pathway.....	27
2.9.4 Type IV secretion pathway.....	28
2.9.5 Type V secretion pathway .....	29

<b>CHAPTER THREE</b> .....	<b>31</b>
<b>MATERIALS AND METHODS</b> .....	<b>31</b>
3.1 MATERIALS .....	31
3.1.1 Bacterial strains .....	31
3.1.2 Cloning and expression vectors.....	31
3.1.3 Oligonucleotide primers .....	31
3.1.4 Chemicals, reagents and media .....	31
3.1.5 Preparation of common media.....	31
3.1.5.1 Mueller-Hinton (MH) broth.....	31
3.1.5.2 Mueller-Hinton (MH) agar .....	34
3.1.5.3 Luria-Bertani (LB) broth .....	34
3.1.5.4 Luria-Bertani (LB) agar .....	34
3.1.5.5 LB agar with ampicilin .....	34
3.1.5.6 LB agar with kanamycin.....	35
3.1.5.7 Chocolate agar .....	35
3.1.5.8 CCDA agar .....	35
3.1.6 Preparation of common buffers and reagents.....	36
3.1.6.1 Tris-EDTA Buffer (TE), pH 8.0.....	36
3.1.6.2 Phosphate buffered saline (PBS), pH 7.2 .....	36
3.1.6.3 PBS-Tween 20 (PBS-T), 0.05% (v/v) .....	36
3.1.6.4 Tris buffered saline (TBS).....	36
3.1.6.5 TBS-Tween 20 (TBS-T), 0.05% (v/v).....	36
3.1.6.6 TBS/Tween 20/Triton-X (TBS-TT) (w/v/v).....	37
3.1.6.7 Lysis buffer.....	37
3.1.6.8 Elution buffer (EB), pH 6.8 .....	37
3.1.6.9 NaOH solution (3M).....	37
3.1.6.10 HCl solution (1M).....	38
3.1.6.11 EDTA solution, 0.5 M (pH 8.0).....	38
3.1.6.12 Ampicillin stock solution (100 mg/ml).....	38
3.1.6.13 Kanamycin stock solution (30 mg/ml) .....	38
3.1.6.14 Chloramphenicol stock solution (34 mg/ml) .....	38
3.1.6.15 Magnesium chloride (MgCl <sub>2</sub> ), 100 mM .....	39
3.1.6.16 Calcium chloride (CaCl <sub>2</sub> ), 100 mM.....	39
3.1.6.17 X-gal, 20 mg/ml (w/v) .....	39
3.1.6.18 IPTG, 800 mM.....	39
3.1.6.19 Ethanol (70%).....	39
3.1.6.20 Restriction enzyme .....	40
3.1.7 Preparation of reagents for agarose gel electrophoresis .....	41
3.1.7.1 Tris-Acetate-EDTA (TAE) buffer (50 × stock).....	41
3.1.7.2 Ethidium bromide solution (EtBr), 10 mg/ml (w/v).....	41
3.1.7.3 Agarose gel loading dye .....	41
3.1.8 Preparation of reagents for SDS-PAGE .....	41
3.1.8.1 Resolving gel buffer, pH 9.3.....	41
3.1.8.2 Stacking gel buffer, pH 6.8.....	42
3.1.8.3 2X Sample buffer.....	42
3.1.8.4 Ammonium persulfate (AP), 20% (w/v) .....	42
3.1.8.5 Running buffer.....	43
3.1.8.6 Coomassie blue stain .....	43
3.1.8.7 Coomassie destaining solution .....	43

3.1.9	Preparation of reagents for immunodetection .....	44
3.1.9.1	Western blot transfer buffer .....	44
3.1.9.2	Blocking stock solution, 10% (w/v) .....	44
3.1.9.3	Staining using chemiluminescence substrate.....	44
3.1.9.4	Ponceau S stain .....	44
3.1.10	Preparation of reagents for histidine-tagged protein purification .....	45
3.1.10.1	Protease inhibitor cocktail (0.37 mg/ml) .....	45
3.1.10.2	Lysozyme stock (10 mg/ml) (w/v) .....	45
3.1.10.3	DNAse 1 (2500 µg/ml) (w/v).....	45
3.1.10.4	10 mM lysis buffer (containing 300 mM NaCl) .....	45
3.1.10.5	20 mM wash buffer (containing 300 mM NaCl) .....	46
3.1.10.6	30 mM wash buffer (containing 300 mM NaCl) .....	46
3.1.10.7	40 mM wash buffer (containing 300 mM NaCl) .....	46
3.1.10.8	10 mM lysis buffer (containing 500 mM NaCl) .....	46
3.1.10.9	20 mM wash buffer (containing 500 mM NaCl) .....	47
3.1.10.10	30 mM wash buffer (containing 500 mM NaCl) .....	47
3.1.10.11	40 mM wash buffer (containing 500 mM NaCl) .....	47
3.1.10.12	Elution buffer (containing 500 mM NaCl) .....	47
3.2	METHODS.....	48
3.2.1	Bacterial strains and growth conditions .....	48
3.2.2	<i>In silico</i> analysis of secreted protein sequences. ....	48
3.2.3	Extraction of the genomic DNA.....	49
3.2.4	Quantification of nucleic acids.....	50
3.2.5	Amplification of <i>CjS8</i> gene using PCR.....	50
3.2.6	DNA agarose gel electrophoresis .....	51
3.2.7	Purification of PCR products .....	52
3.2.8	Preparation of competent cells .....	53
3.2.9	Transformation of plasmid into <i>E. coli</i> TOP10F' competent cells.....	53
3.2.10	Transformation of plasmid into <i>E. coli</i> BL21(DE3) and JM109 .....	54
3.2.11	Long-term storage of recombinant plasmid.....	55
3.2.12	Ligation.....	55
3.2.13	Dephosphorylation of restriction digested DNA .....	56
3.2.14	Protein analysis by SDS-PAGE.....	56
3.2.15	Simply blue safe stain .....	58
3.2.16	Silver stain .....	58
3.2.17	Electrophoretic transfer of proteins to membrane (Western blotting) .....	58
3.2.18	Immunoblotting .....	59
3.2.19	Immunoreactivity study of the fusion proteins by Western blot analysis .....	60
3.2.20	Membrane developing using peroxidase detection method....	60
3.2.21	Oligonucleotide designs of <i>CjS8</i> gene .....	60
3.2.22	Amplification the full length of <i>CjS8</i> gene .....	61
3.2.23	Extraction of pQE70 vector .....	61
3.2.24	Restriction digestion of pQE70 vector .....	61
3.2.25	Gel- purification of digested vector .....	62

3.2.26	Cloning of <i>CjS8</i> gene into pQE70 vector .....	62
3.2.27	Cloning of C-terminal ( <i>CjS8</i> gene) into pCRT7-NT TOPO... ..	63
3.2.28	Expression of C-terminal of <i>CjS8</i> -pCRT7-NT TOPO.....	63
3.2.29	Induction of C-terminal ( <i>CjS8</i> ) protein (pCRT7-NT TOPO) in BL21(DE3)pLysS.....	63
3.2.30	Cloning of C-terminal of <i>CjS8</i> gene in pQE30.....	63
3.2.31	Expression of C-terminal of <i>CjS8</i> -pQE30 .....	64
3.2.32	Induction of C-terminal- <i>CjS8</i> protein (pQE30) in BL21(DE3)pLysS.....	64
3.2.33	Induction of recombinant C-terminal- <i>CjS8</i> protein (pCRT7- NT TOPO) in BL21-CodonPlus-RIL .....	64
3.2.34	Small scale of <i>CjS8</i> protein expression .....	65
3.2.35	Purification of a recombinant C-Terminal protein of <i>CjS8</i> (pCRT7-NT TOPO) in <i>E. coli</i> BL21-CodonPlus-RIL .....	65
3.2.36	Large scale protein expression.....	65
3.2.37	Purification of <i>CjS8</i> protein using gel electro-elution.....	66
3.2.38	Confirmation the purified protein.....	66
3.2.39	Determination of protein solubility .....	66
3.2.40	Production of polyclonal antibody of <i>CjS8</i> protein in rabbit (Ab_ <i>CjS8</i> ) .....	67
3.2.41	Pre-adsorbtion of antibodies .....	68
3.2.42	Construction of <i>C. jejuni</i> NCTC 11168 <i>CjS8</i> gene mutant ( <i>C. jejuni</i> Δ <i>CjS8</i> ).....	68
3.2.43	Amplification, Cloning of the flanking <i>CjS8</i> regions and inverse PCR.....	68
3.2.44	Kanamycin cassette and inverse PCR product .....	69
3.2.45	Transformation of the mutagenesis construct into <i>C.jejuni</i> NCTC 11168 .....	70
3.2.46	Strain survey of clinical isolates of <i>C. jejuni</i> .....	70
3.2.47	Characterization of <i>CjS8</i> as secreted protein and sub- cellular protein fractions localisation .....	71
3.2.48	Involvement of <i>CjS8</i> on adhesion and invasion .....	71
3.2.49	Involvement of <i>CjS8</i> in the formation of biofilm .....	73
3.2.50	Involvement of <i>CjS8</i> in the <i>C. jejuni</i> motility .....	74
3.2.51	Cell culture technique .....	74
<b>CHAPTER FOUR.....</b>		<b>76</b>
<b>RESULTS .....</b>		<b>76</b>
4.1	<i>In silico</i> ANALYSIS FOR THE IDENTIFICATION OF SECRETED PROTEIN SEQUENCE FROM OTHERS PATHOGENIC BACTERIA STRAINS .....	76
4.2	<i>In silico</i> ANALYSIS FOR THE IDENTIFICATION OF SECRETED PROTEINS IN THE <i>C. jejuni</i> NCTC 11168 GENOME .....	76
4.2.1	<i>In silico</i> analysis on C- and N-terminal for the identification of secreted proteins characteristics .....	79
4.3	CLONING AND EXPRESSION OF <i>CjS8</i> GENE IN <i>E. coli</i> .....	82
4.3.1	Cloning of C-terminal <i>CjS8</i> gene in <i>E. coli</i> .....	86
4.3.1.1	Cloning of PCR products into pCRT7/NT-TOPO.....	86
4.3.1.2	Cloning of PCR products into pQE-30 vector.....	89

4.3.2 Cloning of CjS8 gene without the signal peptide (CjS8 wsp) in <i>E. coli</i> .....	94
4.3.3 Protein Expression of C-terminal CjS8 and CjS8 wsp.....	101
4.3.3.1 Small-scale expression.....	101
4.3.3.2 Induction of the target proteins at different time point.....	101
4.3.4 Purification of C-terminal CjS8 recombinant protein in <i>E. coli</i> .....	108
4.3.5 Production of CjS8 specific polyclonal antibody (Ab_CjS8).....	112
4.4 MUTAGENESIS OF CJS8.....	114
4.4.1 Oligonucleotide primers.....	114
4.4.2 Cloning of flanking CjS8 gene in pGEM-T vector.....	114
4.4.3 Deletion of <i>CjS8</i> gene in the pGEM-T construct using inverse PCR mutagenesis.....	120
4.4.4 Insertion of Kanamycin resistant gene ( <i>Kan<sup>r</sup></i> ) into pGEM-T (without CjS8).....	120
4.4.5 Analysis of pGEM-T-without <i>CjS8-Kan<sup>r</sup></i> .....	124
4.4.5.1 Restriction endonuclease digestion.....	124
4.4.5.2 PCR screening of the recombinant plasmids.....	126
4.4.6 Transformation of pGEM-T-without <i>CjS8-Kan<sup>r</sup></i> construct into the <i>C. jejuni</i> NCTC 11168.....	128
4.4.7 Confirmation of CjS8 gene expression in <i>C. jejuni</i> NCTC 11168 and <i>C. jejuni</i> ΔCjS8 using immunoblot analysis.....	132
4.5 CHARACTERISATION OF CJS8 AS A PATHOGENIC DETERMINANT IN <i>C. jejuni</i> .....	134
4.5.1 Sub-cellular localisation of CjS8.....	134
4.5.2 Survey on clinical isolate of <i>C. jejuni</i> for the presence of CjS8 gene.....	138
4.5.3 Involvement of <i>CjS8</i> gene on adhesion and invasion of human epithelial colorectal adenocarcinoma (Caco-2) cells.....	144
4.5.4 Involvement of <i>CjS8</i> gene on the formation of biofilm.....	144
4.5.5 Involvement of <i>CjS8</i> gene on the motility of <i>C. jejuni</i> .....	148
<b>CHAPTER FIVE.....</b>	<b>150</b>
<b>DISCUSSION AND CONCLUSION.....</b>	<b>150</b>
5.1 BIOINFORMATIC ANALYSIS AND CLONING OF CjS8.....	150
5.2 PRODUCTION OF POLYCLONAL ANTIBODY AB_CJS8.....	158
5.3 DEVELOPMENT OF CJS8 MUTANT STRAIN ( <i>C. jejuni</i> ΔCJS8) OF <i>C. jejuni</i> .....	163
5.4 SUMMARY AND RECOMMENDATION FOR FUTURE RESEARCH.....	164
5.4.1 Future directions.....	166
<b>REFERENCES.....</b>	<b>168</b>
<b>APPENDICES.....</b>	<b>198</b>
APPENDIX A – Nucleotide sequence of CjS8 (NC_002163.1).....	198
APPENDIX B – Selected secreted protein sequence from others pathogenic bacteria.....	200
APPENDIX C – Selected <i>C. jejuni</i> sequences obtained from the <i>in silico</i> analysis.....	203
APPENDIX D – Phylogenetic and homology analyses of CjS8 with selected secreted proteins from others pathogenic bacteria.....	206
<b>PUBLICATIONS DURING THE PERIOD OF STUDY.....</b>	<b>207</b>

## LIST OF TABLES

<u>Table No.</u>		<u>Page No.</u>
Table 3.1	Bacteria strain used in this study	37
Table 3.2	Plasmid used in this study	38
Table 3.3	Primers used in this study (Sequence directions: 5`-3`)	38
Table 4.1	Selected secreted protein sequence from others pathogenic bacteria stains. Full sequence of all the proteins are listed in Appendix C	81
Table 4.2	Selected protein sequence of <i>C. jejuni</i> from the NCTC 11168 genome database obtained through <i>in silico</i> protein-protein sequence analysis against known secreted protein of other pathogenic bacteria. Full sequence of all the proteins are listed in Appendix C	82
Table 4.3	<i>In silico</i> analysis for secreted protein characterization on signal peptide cleavage site (▼) at N-terminal and determination of protein motif at C-terminal	84

## LIST OF FIGURES

<u>Figure No.</u>		<u>Page No.</u>
1.1	Flowchart of the study	7
4.4	Bioinformatics analysis on selected pathogenic bacteria and <i>C. jejuni</i> (Cj_) peptide sequences using the SignalP 5.0, Motif Scan (Pfam/PROSITE) and Uniprot PTM protein databases	86
4.5	A). Morphology of Cj8 and B). Protein sequence of CjS8 (GenBank: MPO30128.1)	88
4.6	Complete sequence of <i>CjS8</i> gene (Genebank ID: NC_002163.1). The position <i>CjS8</i> gene in the genome of <i>C. jejuni</i> NCTC 11168 is 1297691-1300816. The nucleotide sequence of <i>CjS8</i> gene was translated into protein sequence using Expasy Translate tool ( <a href="https://web.expasy.org/translate/">https://web.expasy.org/translate/</a> )	89
4.7	PCR product of C-terminal CjS8 gene on 1% DNA agarose gel electrophoresis	91
4.8	Map and sequence characteristics of pCRT7/NT-TOPO vector shows the cloning region	92
4.9	Restiction enzyme analysis of C-terminal CjS8/pCRT7-NT-TOPO vector resolved on 1% DNA agarose gel electrophoresis	94
4.10	Map and sequence characteristics of pQE-30 vector shows the cloning region (MCS- multiple cloning site)	95
4.11	Restiction enzyme analysis of C-terminal CjS8/pQE-30 vector resolved on 1% DNA agarose gel electrophoresis	96
4.12	Figure Map and sequence characteristics of pQE-70 vector shows the cloning region (MCS- multiple cloning site) and selected restriction enzyme sites on pQE-70 and CjS8 <i>wsp</i> sequence genes. The position of enzymes digestion site was identified	99
4.13	Agarose gel electrophoresis analysis of PCR amplification of <i>CjS8 wsp</i> with expected size 3075 bp	100
4.14	Restriction enzyme analysis of pQE-70-CjS8 <i>wsp</i> using <i>Bam</i> HI resolved on 1% DNA agarose gel electrophoresis	101

4.15	PCR screening of pQE-70-CjS8 <i>wsp</i> recombinant resolved on 1% agarose gel electrophoresis. PCR detection using <i>CjS8 wsp</i> primers	102
4.16	Restriction enzyme analysis of pQE-70-CjS8 <i>Swp</i> recombinant cloned resolved on 1% DNA agarose gel electrophoresis	103
4.17	SDS-PAGE analysis of C-terminal CjS8 recombinant protein in pCRT7-NT TOPO induced at different post-induction time point in <i>E. coli</i> BL21(DE3) pLys	105
4.18	SDS-PAGE analysis of C-terminal CjS8 recombinant protein in pCRT7-NT TOPO induced at different post-induction time point in <i>E. coli</i> BL21(DE3) pLys	106
4.19	SDS-PAGE analysis of C-terminal CjS8 recombinant protein in pCRT7-NT TOPO induced at different post-induction time point in <i>E. coli</i> BL21(DE3) pLys.	107
4.20	SDS-PAGE analysis of C-terminal CjS8 recombinant protein in pCRT7-NT TOPO induced at different post-induction time point in <i>E. coli</i> BL21-CodonPlus- RIL	109
4.21	Western blot analysis of C-terminal CjS8 recombinant protein expressed in supernatant and pellet. The expression of C-terminal CjS8 in BL21-CodonPlus-RIL competent cells cloned in pCRT7-NT TOPO vector	110
4.22	Ni-NTA chromatography purification of C-terminal CjS8 after regenerating the column (double loaded) of the cleared lysate (A): SDS-PAGE analysis. (B): Western blot analysis was probed using anti penta-His as primary antibody and developed by using chloronaphthol. (C): Western Blot analysis was probed using anti penta-His as primary antibody and developed by using ECL western blot analysis system	112
4.23	SDS-PAGE stained gel for protein purification of C-terminal CjS8 (34 kDa)	113
4.24	Western blot analysis of C-terminal CjS8 purified protein. The protein was probed using anti penta His as primary antibody and Goat anti Mouse-HRP as the secondary antibody and developed using ECL western blot analysis system	114


4.25	Western blot analysis to examine the Ab_CjS8 against purified C-terminal CjS8 protein. The membrane was developed using ECL western blot analysis system	116
4.26	The position of the PCR primers (A) and schematic diagram (B) of <i>CjS8</i> gene for the mutagenesis experiment	118
4.27	(A) Map and sequence characteristics of pGEM-T Easy Vector shows the cloning region (MCS- multiple cloning site) and <i>NotI</i> restriction enzyme sites on pGEM-T/CjS8 (position 43 and 5217). (B) The restriction enzymes site (pGEM-T/CjS8 with flanking DNA) were generated using NEBcutter V2.0 ( <a href="http://nc2.neb.com/NEBcutter2/index.php">http://nc2.neb.com/NEBcutter2/index.php</a> ).	119
4.28	PCR product of 5.2 kb <i>CjS8</i> with flanking DNA region	120
4.29	Restriction enzyme analysis of <i>NotI</i> on constructed clone pGEM-T/CjS8 with flanking DNA	121
4.30	PCR product of amplification using inverse PCR primers specific for deletion of <i>CjS8</i> gene in the pGEM-T vector	123
4.31	Gel electrophoresis showing the restriction digestion by <i>BamHI</i> of pJMK30 construct as expected size	124
4.32	Restriction enzyme analysis ( <i>BamHI</i> ) of the pGEM-T-without <i>S8-Kan<sup>r</sup></i> construct on 1% agarose gel electrophoresis	126
4.33	PCR screening of pGEM-T-without <i>CjS8-Kan<sup>r</sup></i> recombinant on 1% agarose gel electrophoresis	128
4.34	PCR screening of <i>CjS8</i> gene in the <i>C. jejuni</i> Δ <i>CjS8</i> resolved on 1% agarose gel electrophoresis	130
4.35	PCR screening of Kanamycin resistance gene ( <i>Kan<sup>r</sup></i> ) in <i>C. jejuni</i> Δ <i>CjS8</i> resolved on 1% agarose gel electrophoresis	131
4.36	PCR screening of <i>CjS8</i> gene with 1 kb flanking DNA regions (N- and C-terminal of <i>CjS8</i> gene) in <i>C. jejuni</i> Δ <i>CjS8</i> resolved on 1% agarose gel electrophoresis	132
4.37A	Western blot analysis for the confirmation of <i>CjS8</i> gene expression in <i>C. jejuni</i> NCTC 11168 and <i>C. jejuni</i> Δ <i>CjS8</i> . Equal amount of proteins loaded. Probed using polyclonal antibody Ab_CjS8 (Section 3.2.40) developed by using ECL western blot analysis system detection reagent	134

4.37B	Western blot analysis for the confirmation of <i>CjS8</i> gene expression in <i>C. jejuni</i> NCTC 11168 and <i>C. jejuni</i> Δ <i>CjS8</i>	134
4.38	(A) and (B). SDS-PAGE analysis on secreted proteins and cellular localization of <i>CjS8</i> protein in <i>C. jejuni</i> NCTC 11168 and <i>C. jejuni</i> Δ <i>CjS8</i>	137
4.39	(A) and (B). SDS-PAGE analysis on secreted proteins and cellular localization of <i>CjS8</i> protein in <i>C. jejuni</i> NCTC 11168 and <i>C. jejuni</i> Δ <i>CjS8</i>	138
4.40	PCR product for survey on clinical isolate of <i>Campylocater</i> spp. for the presence of <i>CjS8</i> gene	140
4.41	Western blot analysis of Ab_ <i>CjS8</i> antibody probed against whole-cell protein lysate from <i>Campylobacter</i> spp.	141
4.42	PCR product for survey on clinical isolate of <i>Campylocater</i> spp for the presence of C-terminal <i>CjS8</i> gene	142
4.43	Western blot analysis of Ab_ <i>CjS8</i> antibody probed against whole-cell protein lysate from <i>Campylobacter</i> spp.	143
4.44	PCR product for survey on clinical isolate of <i>Campylocater</i> spp. for the presence of <i>CjS8</i> gene	144
4.45	Influneece of <i>CjS8</i> on adhesion to and invasion of Caco-2 cells by <i>C. jejuni</i> NCTC 11168 and <i>C. jejuni</i> Δ <i>CjS8</i>	146
4.46	Biofilm formation in <i>C. jejuni</i> NCTC 11168 and <i>C. jejuni</i> Δ <i>CjS8</i> at 24, 48 and 72 hours	147
4.47	Biofilm formation in <i>C. jejuni</i> NCTC 11168 and <i>C. jejuni</i> Δ <i>CjS8</i> at 24, 48 and 72 hours	148
4.48	Influence of <i>CjS8</i> gene on motility of <i>C. jejuni</i> NCTC 11168 and <i>C. jejuni</i> Δ <i>CjS8</i> at 24, 48, 96, 120 and 144 hours. <i>C. jejuni</i> 81116 <i>flaAB</i> <sup>-</sup> mutant (control for non-motile) and <i>C. jejuni</i> 81116 (control for highly motile)	150

## ABBREVIATIONS

Amp	Ampicillin
ATP	Adenosine tri-phosphate
bp	Base pair
C	Celcius
CCDA	Cefoperazone charcoal de-oxy chocolate agar
cDNA	Complimentary DNA
CFU	Colony forming unit
CIAP	Calf Intestinal Alkaline Phosphatase
CjS8	<i>C. jejuni</i> S8 family serine peptidase
<i>C. jejuni</i>	<i>Campylobacter jejuni</i>
<i>C. coli</i>	<i>Campylobacter coli</i>
<i>C. jejuni</i> ΔCjS8	<i>C. jejuni</i> mutant (S8 family serine peptidase)
CjS8 wsp	<i>C. jejuni</i> S8 family serine peptidase without signal peptide
CO <sub>2</sub>	Carbon dioxide
dATP	De-Oxy Adenosine Tri Phosphate
dH <sub>2</sub> O	Distilled water
DMEM	Dulbecco's Modified Eagle's Medium
DNA	Deoxy ribonucleic acid
EB	Envelope buffer
<i>E. coli</i>	<i>Escherichia coli</i>
FCS	Fetal Calf Serum
g	Gram
GBS	Guillain-Barré Syndrome
h	Hour
His	Histidine
HRP	Horse radish peroxidase
Kan	Kanamycin
kb	Kilo base pair
kDa	Kilo Dalton
LB	Luria-Bertani
mÅ	milli Amp
MH	Mueller-Hinton
min	Minute

Ni-NTA	Nickel-nitriotriacetic acid
OD	Optical density
ORF	Open reading frame
PBS	Phosphate buffered saline
PCR	Polymerase chain reaction
RNA	Ribonucleic acid
rpm	revolutions per minute
RT	Room temperature
SDS-PAGE	Sodium dodecyl sulphate-polyacrylamide gel electrophoresis
TAE	Tris-acetate EDTA
UV	Ultra-violet
V	Volts
µg	Microgram
µl	Microliter
µM	micro Molar



# CHAPTER ONE

## INTRODUCTION

### 1.1 BACKGROUND OF THE STUDY

The description of this study must begin from the discovery of the cell surface protein that was found to be specific for *Campylobacter jejuni* (Salleh, 1994) (<https://elib.usm.my-Thesis PhD>). Genomic and protein analysis using bioinformatics tools on that protein reveals the presence of signal peptide at N-terminal of its peptide sequence, thus signifies this protein is highly potential acted as secreted protein (Abdul Wahab, 2000) (<https://elib.usm.my-Thesis MSc>). The sequence of this protein was deposited in Uniprot database named as “*Campylobacter jejuni* strain USM1-putative periplasmic protein” and could be retrieved at <https://www.uniprot.org/citations/-4646025062340752923>. The size of the protein was calculated as 24.1 kDa containing the YCE gene domain (UniProtKB - Q79JB5) identified as putative periplasmic protein at the C-terminal. The gene was also known as “*Campylobacter jejuni* hypothetical protein Cj0419” in the Uniprot database which could be retrieved at <https://www.uniprot.org/citations/SIP78F205116FDE3799>. Based on the information as described above, principally, the whole main ideas and the initial experimental designed of this study was to construct the 24.1 kDa gene (Cj0419) *C. jejuni* mutant for the characterization of its gene function and how this gene is involved in the pathogenesis of *C. jejuni*. The strategies for identifying secreted protein in established genome sequence of *C. jejuni* (NCTC 11168) was based on the protein sequence information of *C. jejuni* secreted protein 24.1 kDa (Abdul Wahab, 2000). Others secreted proteins which were experimentally validated from selected

enterobacteriaceae pathogenic bacteria using proteomic and genome-based computational prediction (Chen *et al.*, 2019).

The secreted protein of an organism represents the proteins that released by all types of cells of living things (Chua *et al.*, 2012). Secretory protein systems are important for many physiological functions of cells or organisms such as sustaining cell-cell communication, proliferation, metabolism (Zhang *et al.*, 2014), immunomodulation (Toapanta *et al.*, 2018) and invasion into host cells to cause the disease (Jang *et al.*, 2020). Markedly, many secreted proteins have been identified as important biomarkers and therapeutic targets (Walker *et al.*, 2017). Therefore, understanding the biological functional of bacterial secreted proteins have great potential to provide a valuable resource for diagnosis, prognosis, and treatment of bacterial diseases (Brown *et al.*, 2013).

*C. jejuni* infections are a major cause of diarrheal disease worldwide. The incidence and prevalence of campylobacteriosis have increased in both developed and developing countries over the last 10 years (Kaakoush *et al.*, 2015). They are the leading cause of foodborne illness, with 56,729 cases reported of which almost 96.37 infection over 100,000 populations at England in 2017 (Public Health England, 2017) and among the main confirmed aetiology of bacterial foodborne illness in USA in 2017, infected approximately of 1.5 million people (CDC, 2017). In one of the study in Malaysia it was reported *C. jejuni* was the most predominantly isolated species (69.5%) from broiler chickens and chicken meat (Sinulingga *et al.*, 2020).

## 1.2 PROBLEM STATEMENT AND SIGNIFICANCE OF THE STUDY

Pathogenic bacteria have the ability to colonized human cells to cause disease. Indeed, bacteria successfully evolved to acclimate at different environments for survival. In addition, bacterial pathogens have to overcome host innate and adaptive immune system, including the microbiome especially in the gut. The secreted proteins of pathogenic bacteria is believed to be the evasion system against the host defence systems. The protein secretion systems of pathogenic bacteria are machineries used to secrete proteins in the extracellular medium, or directly into the targeted cell. Several secretion systems have been described in the scientific literature. Gram-negative bacteria have evolved eight secretion systems and they are made of proteins. The outer membrane proteins (OMPs) or surface protein of the pathogenic bacteria are the main protein that will be translocated through the inner membrane and then inserted in the outer membrane, for full virulence.

*C. jejuni* contamination is common on chicken in grocery stores (Kramer *et al.*, 2000; Guirin *et al.*, 2019; Walker *et al.*, 2019; Sinulingga *et al.*, 2020), even on the outside of packages (Burgess *et al.*, 2005; Chen *et al.*, 2018), as it is a commensal in birds (Skirrow & Blazer, 1995) allowing it to quickly spread unnoticed through a flock (Hermanns *et al.*, 2011) potentially accelerated by flies (Hald *et al.*, 2008). High throughput carcass processing exaggerates the problem by spreading contamination between birds. Chicken 'liquor', the juices evacuated from thawed chicken carcasses, has been found to be an efficient protectant for *C. jejuni* (Coates *et al.*, 1987). Chicken carcasses have as high as 5 logs of *C. jejuni* per bird with prevalences of 77% to 94% of carcasses contaminated (Kramer *et al.*, 2000; Garin *et al.*, 2012; Nohra *et al.*, 2018; Guirin *et al.*, 2019). All of the reported cases above clearly portrayed the important of

*C. jejuni* to human health. Human could be easily infected by the *C. jejuni*, thus it is important to understand on the pathogenesis of *C. jejuni* to human host.

In this study we systematically explored the *C. jejuni* genome information for the identification of secreted protein that involved in the pathogenesis of *C. jejuni* infection. For that purposes the 24.1 kDa surface protein of *C. jejuni* (Salleh, 1994; Abdul Wahab, 2000) and others identified secreted protein from pathogenic enterobacteriaceae bacteria were used in the homology searched using BLAST. This study also will further validated and characterized selected *C. jejuni* secreted proteins using several bioinformatics tools. Furthermore, this study will perform the functional analysis of identified secreted protein of *C. jejuni* through constructing the mutant using inverse PCR mutagenesis. It is believed the outcome of this research provides a valuable resource and crucial data for further elaborating the involvement of secreted protein in the pathogenesis of *C. jejuni* infections.

### **1.3 RESEARCH QUESTIONS**

The present study aims to answer the following questions:

- What are the secreted protein in the *C. jejuni* genome that have the closest homology to the 24.1 kDa and selected secreted protein from the pathogenic bacteria?
- What are the best vector system for cloning and protein expression for the *C. jejuni* secreted protein?
- What is the specificity of recombinant secreted protein of *C. jejuni* polyclonal antibody raised in rabbit?